

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

ACC 4.1-15
1/16/2003

ENHANCED HERBICIDES

CROSS REFERENCE TO RELATED APPLICATIONS

This application is a continuation of application Serial No. 09/692,763, filed on October 19, 2000, which is a division of Serial No. 09/427,476, filed October 26, 1999, now U.S. Patent No. 6,218,336 dated April 17, 2001.

BACKGROUND OF THE INVENTION

The invention relates generally to herbicides, methods of improving existing herbicides and controlling the growth of plant life and more particularly to methods and products involving succinate-based chemicals and other additives which can enhance the activity (effectiveness) of herbicidal compounds. As used herein, herbicidal refers to materials which destroy or inhibit plant growth, such as by desiccation or defoliation, for example, to act as a harvest aid or to control weed growth.

Glyphosate and paraquat are the number 1 and 2 non-selective herbicides used worldwide. Paraquat is extremely toxic and therefore unacceptable for many applications. Glyphosate can be slow acting, commonly requiring 1 to 2 weeks to achieve plant death and is therefore also unsuitable for many herbicide applications.

Other conventionally known herbicides include fatty acids, such as pelargonic acid, a nine carbon fatty acid, and caprylic acid, an eight carbon fatty acid. Scythe, sold by Mycogen/Dow and Liberty, made by AgrEvo are known commercially available herbicides. Pelargonic acid is the active ingredient in SCYTHER and glufosinate-ammonium is the active ingredient in LIBERTY. However, the activity of these products is such that the cost of products such as SCYTHER can be undesirably high and the amount of active ingredients needed in products such as paraquat could lead to undesirable effects.

Accordingly, it is desirable to provide improved herbicides, methods for enhancing the activity of existing herbicides and methods of controlling plant growth in order to overcome inadequacies of the prior art.

SUMMARY OF THE INVENTION

Generally speaking, in accordance with the invention, improved herbicide compositions, methods of making the compositions and methods of controlling plant growth are provided. The subject invention relates to combining fatty acid based and other herbicides with succinic acid, succinic acid derivatives and other additives such as those having essentially no herbicidal activity to increase the activity of the herbicide and provide methods of controlling plant growth by applying a combination of a herbicidal fatty acid with succinic acid and/or succinic acid derivative chemicals and/or other additives. In addition to the use of succinic acid, combining other Krebs cycle acids with herbicides can also provide beneficial effects. Concentrations of herbicides and additives applied to plants in accordance with the invention can include compositions involving about 0.1 to 30% herbicide and additive, preferably 0.5 to 15% herbicide and additive and more preferably 1-8% herbicide and additive. The ratio of herbicide, such as fatty acid herbicide to activity enhancing additive can be from about 1:10 to 20:1, preferably 1:1 to 20:1, more preferably 1:1 to 5:1.

Accordingly, it is an object of the invention to provide improved herbicidal compositions.

Another object of the invention is to provide additives which can enhance the activity of herbicidal compositions.

Yet another object of the invention is to improve the safety of herbicides.

A further object of the invention is to provide improved methods of plant control.

The invention accordingly comprises the several steps and the relation of one or more of such steps with respect to each of the others, and the compositions possessing the characteristics, properties and the relation of constituents useful to effect such steps, which will be exemplified in the compositions hereinafter described, all as exemplified in the following detailed disclosure, and the scope of the invention will be indicated in the claims.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The invention is directed to improved herbicidal compositions. Herbicides, such as those including fatty acids as the active ingredient can be formulated at low fatty acid concentration and have relatively low herbicidal activity or at higher fatty acid concentration and have enhanced activity. However, it has been found that certain additives, such as organic acids and their derivatives, which may have substantially no herbicidal activity can enhance the herbicidal activity of herbicidal fatty acids and other herbicides and provide herbicides with either enhanced activity or reduced fatty acid concentrations. Preferred additives include those organic acids which are part of the Krebs cycle and in particular, succinic acid as well as succinic acid derivatives. Thus, substantially safe non-herbicidal additives are combined with G.R.A.S. (generally

recognized as safe by the FDA) herbicidal compounds such as fatty acids, and the result is a generally safe herbicide with enhanced activity.

The following examples demonstrate the synergistic relationship between additives (such as succinic acid) and fatty acid herbicides such as caprylic acid, pelargonic acid and others. Other synergistic relationships between organic acids (for example, citric acid, tartaric acid, malic acid and lactic acid) with caprylic acid as the fatty acid were also exhibited. In general, with the exception of tartaric acid, there was a general lack of correspondence between the acidity of the organic acid and the degree of synergy of the organic acid with caprylic acid. It was determined that tartaric acid (e.g. L-tartaric acid) exhibited particularly high performance enhancement of caprylic acid across a wide variety of plant types.

A synergistic relationship between succinic acid and sodium salicylate was also demonstrated where the effect of succinic acid alone on the plants treated was negligible.

Accordingly, it has been determined that combining certain organic acids and compounds having a significant herbicidal effect, such as pelargonic, caprylic, caproic, capric and oleic acid, and also such acids as acetic, butyric, valeric, hexanoic and heptanoic acid and compounds such as sodium salicylate, glyphosate (in Round Up) or glufosinate-ammonium with other organic acids and additives including those having substantially no herbicidal effect could enhance herbicidal activity and reduce costs, environmentally undesirable effects or be otherwise more convenient to use.

Effective additives include succinic acid and succinic acid derivatives such as dimethyl succinic acid, calcium succinate, magnesium succinate, diammonium succinate and ammonium succinate as well as certain other organic acids, such as tartaric acid, citric acid, malic acid, lactic acid, adipic acid and plant oils such as limonene and pine oil, especially Unipine (a pine oil derivative available from Busche, Boake & Allen, Inc.), as well as other additives including ammonium sulfate, ammonium tartrate, ammonium chloride and sodium salicylate.

In order to confirm that additives in accordance with the invention provided an enhanced herbicidal effect, a number of experiments were performed in which only fatty acids or other herbicides were applied to plants, the additives alone were applied and the fatty acids plus the additives were applied. The herbicides were applied in the "spray to drip" amount or at a calibrated 20-60 gallons/acre. "Spray to drip" is an uncalibrated application of fluid to foliage, where sufficient spray volume is used to sufficiently saturate the foliage surface until excess fluid begins to drip from the foliage. In general, about 5 to 200 gallons/acre, preferably 20-100 gallons/acre can be effective.

Fatty acids above 6-7 carbon atoms tend to be relatively insoluble in water. Caprylic and pelargonic acids are 8 and 9 carbon acids respectively and require a solvent, such as acetone or an emulsifier to help prevent separation between aqueous and lipid phases. Herbicides can be provided in concentrated form and then diluted at the point of use.

Aspects and embodiments of the invention will be described more clearly with reference to the following examples, which are intended to be interpreted as exemplary, and not in a limiting sense.

Example 1

Potato field trial: two varieties (Snowden and Russet Burbank)
Comparison of Desiccate II with experimental formulations

Average injury rating* & (%vine injury) (days after 1" application)

Treatment:	Snowden		Russet Burbank	
	Day 5	Day 8	Day 5	Day 8
1 Desiccate II (2.175 quart/acre)	0 (0)	30 (10)	50 (40)	60 (70)
2 4% pelargonic + 0.5% succinic acid	90 (70)	100 (94)	97 (90)	98 (95)
3. 4% caprylic/capric + 0.5% succinic	90 (70)	96 (92)	95 (90)	98 (97)
4 4% caprylic/capric + 0.5% succinic + 1% sodium salicylate	90 (80)	100 (98)	95 (90)	96 (92)
5 4% sodium salicylate + 0.5% succinic	25 (0)	40 (10)	70 (60)	98 (92)
6 6% caprylic/capric + 0.5% succinic	95 (75)	100 (90)	80 (70)	98 (94)
				Day 15
				100 (96)
				100 (100)
				100 (100)
				100 (100)
				100 (100)
				100 (100)

* Injury rating based on scale of 1-5, where 5 = complete desiccation of all plants/plot
 Second application 5 days after 1" application; each at 87 G/a (45 psi)
 Treatment #1 contained 0.125% Wilfarm Crop Oil Concentrate
 Treatments #2-4 and 6 contained 0.83% Emisorb 6900, 0.43% mineral oil and 1% Hasten
 Treatment #5 contained 0.3% Sylgard 309
 Summary: all experimental treatments (#2-6) better than treatment #1 (Desiccate II)

Example 2

Potato field trial: 2 applications of selected herbicides, including Desiccate II

Treatment	Injury rating*, average per plot				
	Day 1	Day 2	Day 3	Day 4	Day 5
1 5% pelargonic	4.07	4	4.13	4.43	4.43
2 5% pelargonic + 1% di-ammonium succinate	4.23	4.23	4.47	4.57	4.57
3 5% caprylic/capric + 1% di-ammonium succinate	3.87	3.9	4.03	4.4	4.4
4 Desiccate II (1.5 quarts/acre)	1	1.5	1.67	3	3

* Injury rating, based on a scale of 1-5, where 5= complete desiccation of all plants/plot
 1% Hasten in treatments 1, 2 and 3. For treatment 4: 0.125% Wilfarm Crop Oil Concentrate.

Treatments 1, 2 and 3 also had 0.86% Emsorb 6900 and 0.43% mineral oil
 Plot size: 3X6 feet. Each treatment group had 3 replicates

40-42 G/acre, with <45 psi for 1st application and 45 psi for 2nd application
 2nd treatment applied 2 days after 1st treatment

Variety: Katahdin (thick canopy), with active growth (irrigated)

Summary: 1) treatments 1-3 superior to treatment 4 (Desiccate II),
 2) pelargonic acid enhanced by di-ammonium succinate

Example 3

Dry bean (navy) field trial: single application of selected herbicides, including Desiccate II

Treatment	Injury rating*, average per plot					% desiccation, day 14		
	Day 2	Day 4	Day 5	Day 6	Day 14	foliage	vines	pod
1 Desiccate II (1.5 quarts/acre)	3.375	3.8	4.55	4.68	4.625	90	50	60
2 4% pelargonic + 0.5% succinic acid	4.325	4.35	4.7	4.73	4.763	98	90	97
3 4% caprylic + 0.5% succinic acid	4.413	4.43	4.76	4.83	4.8	98	90	97
4 4% sodium salicylate + 0.5% succinic	4.388	4.38	4.68	4.73	4.775	95	80	97

*Injury rating, based on a scale of 1-5, where 5 = complete desiccation of all plants/plot
 1% Hasten in treatments 2, 3 and 0.3% Sylgard 309 in treatment 4. Treatment 1 had 0.125% Wilfarm Crop Oil Concentrate
 Treatments 2 and 3 also contained 0.86% Emsorb 6900 and 0.43% mineral oil
 Plot size = 3 X 6 feet, with 4 replicates (plots) per treatment group
 40 gallons per acre at 45 psi for one application
 Dry bean variety (navy): Vista
 Field

Summary: Treatments 2-4 were superior to treatment 1 (Desiccate II)

Example 4

Bean (black) trial: single application of selected herbicides, including Desiccate II

Treatment:	Injury rating*, average per plot				Day 9	Day 19	% desiccation, day 19	
	Day 1	Day 3	Day 7	Day 19			foliage	pod.s
1 Desiccate II (1.5 quarts/acre)	1.75	4.35	4.5	4.7	4.88		99	100
2 3% caprylic + 0.5% succinic acid	4.3	4.55	4.55	4.7	4.9		99	100
3 3% pelargonic + 0.5% succinic acid	4.55	4.75	4.78	4.85	5		100	100
4 3% sodium salicylate + 0.5% succinic acid	4.5	4.7	4.7	4.83	5		100	100

*Injury rating, based on a scale of 1-5, where 5 = complete desiccation of all plants/plot
 1% Hastern in treatments 2, 3 and 0.3% Sylgard 309 in treatment 4. Treatment 1 had 0.125% Wilfarm Crop Oil Concentrate
 Treatment 2 and 3 also contained 0.86% Emsorb 6900 and 0.43% mineral oil
 Plot size: 3 X 6 feet, with 2 replicates (plots) per treatment group
 40 gallons per acre at 45 psi for one application
 Dry bean variety: black, T39
 Field

Summary: Treatments 2-4 superior to treatment 1

Example 5

Synergistic interactions of succinic acid and di-ammonium succinate with herbicidal compounds, on navy dry beans

Treatment:	Injury rating*, average per plot				% desiccation (Day 12)			
	Day 1	Day 2	Day 3	Day 4	Day 12	foliage	vines	Pods
1 3% caprylic	3.5	3.93	3.93	4	4.2	50	10	80
2 3% caprylic + 0.5% succinic acid	4.43	4.43	4.6	4.5	4.7	95	70	85
3 3% caprylic + 2% succinic acid	4.5	4.47	4.57	4.5	4.8	99	85	90
4 3% caprylic + 2.52% di-ammonium succinate	4.47	4.43	4.67	4.6	4.8	99	90	99
5 3% sodium salicylate	3.8	4.13	4.57	4.5	4.7	92	70	90
6 3% sodium salicylate + 0.5% succinic acid	3.93	4.27	4.63	4.6	4.8	95	85	92
7 3% sodium salicylate + 2% succinic acid	4.2	4.4	4.67	4.6	4.9	97	80	95
8 3% sodium salicylate + 2.52% di-ammonium succinate	4.4	4.43	4.72	4.6	4.9	97	90	95
9 0.5% succinic acid	0	0	0	0	0	0	0	0
10 2% succinic acid	0	0	0	0	0	0	0	0
11 2.52% di-ammonium succinate	0	0	0	0	0	0	0	0
12 0.5% succinic acid	0	0	0	0	0	0	0	0
13 2% succinic acid	0	0	0	0	0	0	0	0
14 2.52% di-ammonium succinate	0	0	0	0	0	0	0	0

*Injury ratings, based on a scale of 1-5, where 5 = complete desiccation of all plants/plot

Treatments 1-4, 9-11: each in 0.86% Emsorb 6900, 0.43% mineral oil, 1% Hasten

Treatments 5-8, 12-14: each in 0.3% Sylgard 309

Succinic (2%) and di-ammonium succinate (2.52%) are equimolar

Plot size: 3 X 6 feet. Each treatment group had 3 replicates

One application at 40 G/acre, 45 psi

Variety: dry bean (navy Vista)

Field

Summary: both succinic acid and di-ammonium succinate were synergistic with sodium salicylate and caprylic acid

Example 6

Synergistic interactions of succinic acid and di-ammonium succinate with other herbicidal compounds, on dry beans (black)

Treatment:	Injury rating*, average per plot	% desiccation (Day 12)		
	Day 2	Day 12	foliage	stems
1 3% caprylic	4	4.73	80	95
2 3% caprylic + 0.5% succinic acid	4.37	4.87	97	95-98
3 3% caprylic + 2% succinic acid	4.43	4.92	99	98
4 3% caprylic + 2.52% di-ammonium succinate	4.37	4.9	99	95-97
5 3% sodium salicylate	4	4.78	95	90
6 3% sodium salicylate + 0.5% succinic acid	4.23	4.87	99	98
7 3% sodium salicylate + 2% succinic acid	4.07	4.88	99	95
8 3% sodium salicylate + 2.52% di-ammonium succinate	4.37	4.8	99	92
9 0.5% succinic acid	0	0	0	0
10 2% succinic acid	0	0	0	0
11 2.52% di-ammonium succinate	0	0	0	0
12 0.5% succinic acid	0	0	0	0
13 2% succinic acid	0	0	0	0
14 2.52% di-ammonium succinate	0	0	0	0

*Injury rating, based on a scale of 1-5, where 5 = complete desiccation of all plants/plot
 Treatments 1-4, 9-11: each in 0.86% Emsorb 6900, 0.43% mineral oil, 1% Hasten
 Treatments 5-8, 12-14: each in 0.3% Sylgard 309
 Succinic (2%) and di-ammonium succinate (2.52%) are equimolar
 Plot size: 3 X 6 feet. Each treatment group had 3 replicates
 One application at 40 G/acre, 45 psi
 Variety: black beans (dry bean variety T39)
 Summary: both succinic acid and di-ammonium succinate were synergistic with sodium salicylate and caprylic acid

Example 7

Enhancement of caprylic acid/sodium salicylate as an herbicide, by succinic acid

Treatment	Succinic acid (%)	Relative injury rating* (average, based on 2 ratings)		
		potatoes	plants	All
1 0.25% caprylic + 0.25% sodium salicylate	0	3	8.5	
2 0.25% caprylic + 0.25% sodium salicylate	0.5	5	17	
3 0.25% caprylic + 0.25% sodium salicylate	1	6	17.6	
4 0.25% caprylic + 0.25% sodium salicylate	1.5	7.7	19.7	
5 0.25% caprylic + 0.25% sodium salicylate	2	6.8	18.9	
6 0.5% caprylic + 0.5% sodium salicylate	0	4.8	15.1	
7 0.5% caprylic + 0.5% sodium salicylate	0.5	5.3	17.7	
8 0.5% caprylic + 0.5% sodium salicylate	1	5.5	19.7	
9 0.5% caprylic + 0.5% sodium salicylate	1.5	7	21.3	
10 0.5% caprylic + 0.5% sodium salicylate	2	7.5	22.1	

* A rating of 1-5 given (5= complete desiccation of all plants), for each treatment (1-2 plants/pot)
Two independent evaluations (ratings) performed
Helena Kinetix at 0.2% used for all treatments
All formulations in 30% acetone

Spray to drip applications to dry beans, snapbeans and potato plants

Example 8

Effect of 1% succinic acid on herbicidal activity of Liberty and Scythe

Treatment	Total score for all injury ratings*
1 4% Scythe	66
2 2% Scythe	54.2
3 2% Scythe + 1% succinic	56.1
4 2% Scythe + 0.25% caprylic + 0.25% sodium salicylate	56.3
5 2% Scythe + 0.25% caprylic + 0.25% sodium salicylate + 1% succinic	60.3
6 2% Scythe + 0.5% sodium salicylate	47.5
7 Liberty (4oz/gal)	87.8
8 Liberty (2oz/gal)	76.6
9 Liberty (2oz/gal) + 1% succinic	81
10 Liberty (2oz/gal) + 0.25% caprylic + 0.25% sodium salicylate	77
11 Liberty (2oz/gal) + 0.25% caprylic + 0.25% sodium salicylate + 1% succinic	78.4
12 Liberty (2oz/gal) + 0.5% sodium salicylate	75.7
13 Liberty (1oz/gal)	65.9
14 Liberty (1oz/gal) + 1% succinic	70.3
15 Liberty (1oz/gal) + 0.25% caprylic + 0.25% sodium salicylate	67.6
16 Liberty (1oz/gal) + 0.25% caprylic + 0.25% sodium salicylate + 1% succinic	72.1
17 Liberty (1oz/gal) + 0.5% sodium salicylate	70.8

*Injury rating based on 1 to 6, where 6= complete desiccation of plant. Six independent evaluations performed and all scores, based on ratings for nutsedge, snapbeans and Kentucky bluegrass, where given as a total score for each treatment Kinetic at 0.2% used for all treatments. Applications of treatments made via spray to drip. Snapbeans (full bean production), nutsedge (6-8 inches, height) and bluegrass (blade height 3 inches)

1% succinic acid = zero rating. Succinic was synergistic with those combinations given above.

Example 9

Enhancement of RoundUp Ultra's herbicidal activity on snapbeans and potatoes using 1% succinic acid

Treatment	Average injury rating*	
	potatoes	snapbeans
1 RoundUp Ultra, 1.25%	4.48	2.92
2 RoundUp Ultra, 1.25% + 1% succinic acid	4.67	4.78
3 RoundUp Ultra, 0.25%	3.35	2.82
4 RoundUp Ultra, 0.25% + 1% succinic acid	4.37	3.2

*Rating of 1 to 6 = complete desiccation

Average rating based on 6 independent evaluations of injury to plants

Application rate of 1.25% RoundUp Ultra at 40 G/acre = 2 quarts/acre

(0.25% RoundUp Ultra at 40G/acre= 0.4 quarts/acre)

Mature snapbeans (with beans pods) and potatoes at 6-8 inches in height were used

Summary: succinic acid synergistic with RoundUp

Example 10

Succinic acid enhancement of herbicidal activities:
sodium salicylate +/- caprylic acid

Treatment	Injury rating*	
	Spray to drip	Average rating per pot
1 0.75% caprylic + 0.75% sodium salicylate	45.1	2.82
2 0.75% caprylic + 0.75% sodium salicylate + 1% succinic acid	60.3	3.77
3 1% caprylic + 1% sodium salicylate	50.1	3.13
4 1% caprylic + 1% sodium salicylate + 1% succinic acid	61.1	3.82
5 1.5% sodium salicylate	56.2	3.51
6 1.5% sodium salicylate + 1% succinic acid	62.5	3.91

*For each pot, a rating (1 to 5) was given, where 5 was complete desiccation of all plants

Three independent evaluations were done

Test plants for spray to drip application: Kentucky bluegrass, velvetleaf, foxtail, tomato, potato and snapbeans

All spray solutions contained 30% acetone and 0.3% Sylgard 309

Caprylic acid was v/v and succinic acid and sodium salicylate were at wt/v

Summary: Succinic acid enhanced effectiveness

Example 11

Enhancement of herbicide formulations with 1% succinic acid

Treatment		Average Injury rating
1	0.5% caprylic + 0.5% sodium salicylate	31.6
2	0.5% caprylic + 0.5% sodium salicylate	41.4
3	0.5% caprylic + 0.5% sodium salicylate	41.6
4	0.5% caprylic + 0.5% sodium salicylate	39.2
5	0.5% caprylic + 0.5% sodium salicylate	46.8
6	0.5% caprylic + 0.5% sodium salicylate	44.6
7	0.5% caprylic + 0.5% sodium salicylate	43.6
8	0.5% caprylic + 0.5% sodium salicylate	40
	0.5% ammonium chloride	1% succinic acid
	0.5% ammonium chloride	1% succinic acid
	0.5% ammonium chloride	1% succinic acid

*A rating of 1-5, where 5 = complete desiccation of all plants
 Three independent evaluations (ratings) performed
 Helena Kinetic at 0.1% used for all applications
 All formulations in 30% acetone
 Spray to drip applications to snapbeans and potato plants

Example 12

Succinic acid enhancement of herbicidal activities:

sodium salicylate combined with hexanoic, acetic or caprylic/capric acid

Treatment	Injury rating*	
	Spray to drip 44:1	Average rating per pot 3.68
1 2% hexanoic + 1% sodium salicylate	55.8	4.65
2 2% hexanoic + 1% sodium salicylate + 1% succinic		
3 3% acetic + 1% sodium salicylate	47.1	3.93
4 3% acetic + 1% sodium salicylate + 1% succinic	54.4	4.53
5 2% caprylic/capric + 1% sodium salicylate	50	4.17
6 2% caprylic/capric + 1% sodium salicylate + 1% succinic	54.4	4.53
7 2% caprylic/capric + 1% sodium salicylate + 2% succinic	56.7	4.73

*For each pot, a rating (1 to 5) was given, where 5 was complete desiccation of all plants. Two independent evaluations were done.

Test plants for spray to drip application: velvetleaf, corn snapbeans, foxtail and tomatoes. All spray solutions contained 30% acetone and 0.3% Sylgard 309.

Acetic, hexanoic and caprylic/capric acid were v/v and succinic acid was wt/v.

Caprylic/capric (caprylic, capric, caproic and lauric at 58, 40, 1 and 1%, respectively).

Example 13

Succinic acid enhancement of oleic acid/sodium salicylate herbicidal activity

<u>Treatment</u>	<u>Injury rating*</u>	
	<u>Spray</u>	<u>Average</u>
1 2% oleic acid	<u>to drip</u>	<u>rating</u>
2 2% oleic acid + 1% sodium salicylate	<u>per pot</u>	
3 2% oleic acid + 1% sodium salicylate + 1% succinic acid	26.4	2.2
4 2% oleic acid + 1% succinic acid	37.9	3.16
	49.2	4.1
	28.1	2.33

*For each pot, a rating (1 to 5) was given, where 5 was complete desiccation of all plants
Two independent evaluations were done

Test plants for spray to drip: lambsquarter, velvetleaf, foxtail, nutsedge and potatoes
All spray solutions contained 50% acetone and 0.3% Sylgard 309

Oleic acid was v/v and succinic acid and sodium salicylate was wt/v

Summary: Sodium salicylate and succinic acid enhance effectiveness

Example 14

Succinic acid enhancement of oleic acid/sodium salicylate herbicidal activity

Treatment	Total Injury rating*			Average rating per pot
	60 G/a	Spray to drip	Total	
1 2% oleic acid	23.5	22.3	45.8	1.91
2 2% oleic acid + 1% sodium salicylate	52.3	34.7	87	3.63
3 2% oleic acid + 1% sodium salicylate + 1% succinic acid	63.8	37.1	101	4.2

*For each pot, a rating (1 to 5) was given, where 5 was complete desiccation of all plants. Both 60 gallons/acre and spray to drip applications were made.

Two independent evaluations for both 60 G/a and spray to drip applications

Test plants for 60 G/a: lambsquarter, foxtail and velvetleaf

Test plants for spray to drip: snapbeans, corn and tomatoes

All spray solutions contained 50% acetone and 0.3% Sylgard 309

Oleic acid was v/v and succinic acid and sodium salicylate was wt/v

Example 15

Cotton desiccation: comparison of Scythe with formulations containing caprylic/capric, sodium salicylate and succinic acid

Treatment	Average rating* Three evaluations			Average rating for all 3 evaluations
	1	2	3	
1 2% caprylic + 1% sodium salicylate	4	3.5	3.6	3.7
2 2% caprylic + 1% sodium salicylate	3.8	4.1	4	3.97
3 2% caprylic + 1% sodium salicylate	4.2	4.5	4.4	4.4
4 2% Scythe (1.2% active ingredients)	2	2	2	2
5 4% Scythe (2.4% active ingredients)	3.4	3.2	3.3	3.27

* rating based on visual injury rating of 1-5, where 5 = complete desiccation of cotton foliage

Cotton variety = DeltaPine NuCotn 33-b

All treatments (except Scythe) in 30% acetone

Caprylic, capric, caproic and lauric at 58, 40, 1 and 1%, respectively; i.e.

Henkel's Emery 658, used at v/v, in water

Applications at 60 gallons/acre

Example 16

Herbicide activity of caprylic acid/sodium salicylate +/- succinic acid

<u>Treatment</u>	<u>Total injury rating*</u>
1 1% caprylic + 3% sodium salicylate + 0.5% succinic	97.7
2 3% caprylic + 1% sodium salicylate + 0.5% succinic	97.9
3 1% caprylic + 3% sodium salicylate + 1.5% succinic	97.9
4 3% caprylic + 1% sodium salicylate + 1.5% succinic	99.9
5 1% caprylic + 3% sodium salicylate	79.8
6 3% caprylic + 1% sodium salicylate	91.3

*For each pot, a rating (1 to 5) was given, where 5 was complete desiccation of all plants Both 60 gallons/acre and spray to drip applications were made.

Two independent evaluations for both 60 G/a and spray to drip applications

Test plants for 60 G/a: lambsquarter, foxtail and Kentucky bluegrass

Test plants for spray to drip: nutsedge, foxtail and velvetleaf

All spray solutions contained 30% acetone and 0.3% Hasten

Caprylic acid was v/v and succinic acid and sodium salicylate were wt/v

Example 17

Herbicidal activity of caproic acid with other compounds:
succinic acid and sodium salicylate

		<u>Total injury ratings*</u>		
<u>Treatment</u>		<u>60 G/a</u>	<u>Spray to drip</u>	<u>Total score</u>
1	2% caproic	11.2	7.3	18.5
2	2% caproic + 1% succinic	13.8	7.6	21.4
3	2% caproic + 2% succinic	12.7	7.1	19.8
4	3% caproic	13.9	7	20.9
5	3% caproic + 1% succinic	14.1	7.9	22
6	3% caproic + 2% succinic	14.5	8.2	22.7
7	2% caproic + 1% sodium salicylate	13.5	8.7	22.2
8	2% caproic + 1% sodium salicylate + 1% succinic	16.7	9.2	25.9

*For 60 G/a: 4 pots/treatment, testing lambsquarter, pigweed and velvetleaf

*For spray to drip: 2 pots/treatment, testing lambsquarter and foxtail

For each pot (2-20 plants/pot), a rating of 1-5 given, where 5 was complete desiccation of all plants
All solutions included 1% Emsorb 6900 and 0.3% Hasten

Summary: sodium salicylate and succinic acid enhance effectiveness

Example 18

Herbicidal activity of caprylic acid combined with other compounds
(i.e., salts of succinate +/- sodium salicate)

Treatment	Total injury ratings*		
	60 G/a	Spray to drip	Total score
1 2% caprylic			
2 2% caprylic + 1.5% diammonium succinate	61	8.2	69.2
3 2% caprylic + 1.5% potassium succinate	67.9	8.8	76.7
	58	7.6	65.6
4 2% caprylic + 1.5% sodium salicylate			
5 2% caprylic + 1.5% sodium salicylate + 1.5% diammonium succinate	63.7	7.8	71.5
6 2% caprylic + 1.5% sodium salicylate + 1.5% potassium succinate	70	8.6	78.6
	58.9	7.5	66.4

*For 60G/a: 8 total pots/treatment, testing lambsquarter, pigweed, velvetleaf and foxtail

*For spray to drip: 2 total pots/treatment, testing velvetleaf and foxtail

Evaluations made: 2 for plants treated at 60 G/a and 1 evaluation for "spray to drip"

For each pot (2-20 plants/pot), a rating of 1-5 given, where 5 was complete desiccation of all plants

All treatments included 1% Emsorb 6900

Example 19

Efficacy of caprylic with succinic or ammonium succinate, as herbicides in an emulsification system (Emsorb 6900)

		Injury rating*, total score for 2 evaluations		
<u>Treatment</u>	<u>60 G/a</u>	<u>Spray to drip</u>	<u>Total Rating</u>	<u>Average rating per pot</u>
1 2% caprylic	73.6	49.8	123.4	4.11
2 2% caprylic + 1.5% ammonium succinate	76.1	54.5	130.6	4.35
3 2% caprylic + 3% ammonium succinate	76.4	57.7	134.1	4.47
4 2% caprylic + 1.5% succinic acid	80.6	52.4	133	4.43

*For each pot, a rating (1 to 5) was given, where 5 was complete desiccation of all plants
Two independent evaluations done for both 60 G/a and spray to drip
Test plants for 60 G/a: black nightshade, redroot pigweed, lambsquarter, foxtail, velvetleaf and snapbeans
Test plants for spray to drip: redroot pigweed, lambsquarter, velvetleaf and foxtail
All spray solutions contained 1% emsorb (Henkel) and 0.3% Hasten
Caprylic acid was at v/v and succinic acid and ammonium succinate were at wt/v

Example 20

Efficacy of caprylic with succinic or ammonium succinate, as herbicides in an emulsification system (Emsorb 6900)

		Injury rating*, total score for 2 evaluations			
<u>Treatment</u>		60 G/a	Spray to drip	Total Rating	Average rating per pot
1	3% caprylic	142	36.6	178.4	4.05
2	3% caprylic + 0.5% ammonium succinate	145	37.5	182.2	4.14
3	3% caprylic + 1.5% ammonium succinate	145	39.8	185	4.2
4	3% caprylic + 1.5% succinic acid	151	44.6	195.9	4.45

*For each pot, a rating (1 to 5) was given, where 5 was complete desiccation of all plants

Two independent evaluations done for both 60 G/a and spray to drip

Test plants for 60 G/a: black nightshade, redroot pigweed, foxtail, lambsquarter and snapbeans

Test plants for spray to drip: redroot pigweed, velvetleaf, nutsedge and foxtail

All spray solutions contained 1% emsorb (Henkel) and 0.3% Hasten

Caprylic acid was at v/v and succinic acid and ammonium succinate were at wt/v

Example 21

Efficacy of 1.5% succinic acid with oleic, caprylic or pelargonic acid, as herbicides in an emulsification system (Emsorb 6900)

Treatment	Injury rating*, total score for 2 evaluation			Average rating per pot
	60 G/a	Spray to drip	Total Rating	
1 3% oleic	24.2	9.4	33.6	1.2
2 3% oleic + 1.5% succinic	27.3	8.7	36	1.29
3 3% caprylic				
4 3% caprylic + 1.5% succinic	88.1 91.5	35.4 37.5	123.5 129	4.41 4.61
5 3% pelargonic				
6 3% pelargonic + 1.5% succinic	85.3 91.2	37.6 37.4	122.9 128.6	4.39 4.59

*For each pot, a rating (1 to 5) was given, where 5 was complete desiccation of all plants
Two independent evaluations done for both 60 G/a and spray to drip

Test plants for 60 G/a: velvetleaf, black nightshade, redroot pigweed, foxtail and snapbeans

Test plants for spray to drip: black nightshade, velvetleaf, foxtail and snapbeans

All spray solutions contained 1% Emsorb (Henkel) and 0.3% Hasten

Oleic, caprylic and pelargonic at v/v and succinic acid at wt/v

Succinic acid, alone, had a rating of zero. Succinic acid was synergistic with oleic, caprylic and pelargonic acids

Example 22

Efficacy of caprylic + sodium salicylate + succinic acid, as an herbicide in an emulsification system (Emsorb 6900)

Treatment	Injury rating*, total score for 2 evaluation			Average rating per pot
	60 G/a	Spray to drip	Total Rating	
1 2% caprylic + 1.5% succinic	58.5	62	120.5	3.08
2 2% caprylic + 1.5% succinic + 1% sodium salicylate	65	64.1	129.1	3.31
3 3% caprylic + 1.5% succinic	68.7	68.1	136.8	3.51
4 3% caprylic + 1.5% succinic + 1% sodium salicylate	72	71.1	143.1	3.67
5 4% caprylic + 1.5% succinic	71	66.6	137.6	3.53

*For each pot, a rating (1 to 5) was given, where 5 was complete desiccation of all plants
Two independent evaluations done for both 60 G/a and spray to drip
Test plants for 60 G/a: redroot pigweed, velvetleaf, snapbeans, cotton and potatoes
Test plants for spray to drip: black nightshade, lambsquarter, foxtail and snapbeans
All spray solutions contained 1% Emsorb (Henkel) and 0.3% Hasten
Caprylic acid was at v/v and succinic acid and sodium salicylate were at wt/v
Succinic acid, alone, had a rating of zero. Succinic acid was synergistic with oleic, caprylic and pelargonic acids

Example 23

Herbicidal activity: relationships between caprylic acid and succinic acid +/- ammonium sulfate

		Total Injury ratings, combined, for two separate evaluations*		
<u>Treatment</u>	<u>60 G/a</u>	<u>Spray to drip</u>	<u>Total Rating</u>	<u>Average rating/pot</u>
1 4% caprylic acid	60.7	56.1	116.8	4.49
2 4% caprylic acid + 0.5% succinic acid	63.9	57	120.9	4.65
3 4% caprylic acid + 1% succinic acid	61.6	55.6	117.2	4.51
4 4% caprylic acid + 2% succinic acid	59.4	57.4	116.8	4.49
5 4% caprylic acid + 0.5% succinic acid + ammonium sulfate	60.4	56.6	117	4.5
6 4% caprylic acid + 1% succinic acid + ammonium sulfate	64	57.8	121.8	4.68
7 4% caprylic acid + 2% succinic acid + ammonium sulfate	62.1	57.4	119.5	4.6

*For 60 G/a: 7 total pots/treatment, testing redroot pigweed, foxtail and velvetleaf
 For spray to drip: 6 total pots/treatment, testing foxtail, dry beans and pigweed
 For each pot (2-20 plants/pot), a ratings of 1 to 5 given, where 5 was complete desiccation of all plants
 Two independent evaluations (ratings) done for plants treated at both 60 G/a and spray to drip
 All treatments included Emsorb 6900 (Henkel) at 1%
 Caprylic acid and succinic acid, as v/v and wt/v, respectively, in water
 Succinic acid, alone, had a rating of zero. Caprylic acid + succinic acid was synergistic

Example 24

Herbicidal activity: comparison of fatty acids, when combined with succinic acid and sodium salicylate

Treatment	pH, spray solution	60 G/a	Total Injury ratings, combined, for two separate evaluations*		Average rating/pot
			Spray to drip	Total	
1 3% butyric acid + succinic/NaSal	3.8	102.3	52.5	154.8	4.3
2 3% valeric acid + succinic/NaSal	3.9	105	53.8	158.8	4.41
3 3% caproic acid + succinic/NaSal	4.4	104.9	51.9	156.8	4.36
4 3% heptanoic acid + succinic/NaSal	4.5	108.9	54.1	163	4.53
5 3% caprylic acid + succinic/NaSal	4.5	112.4	55.5	167.9	4.67
6 3% pelargonic acid + succinic/NaSal	4.5	111.9	56.6	168.5	4.68
7 3% oleic acid + succinic/NaSal	4.5	105	53.2	158.2	4.39
8 3% caprylic/capric + succinic/NaSal	4.6	114.3	57.3	171.6	4.77

*For 60 G/a: 12 total pots/treatment, testing redroot pigweed, wheat, foxtail and velvetleaf.

For spray to drip: 6 total pots/treatment, testing redroot pigweed, foxtail and velvetleaf

For each pot (2-20 plants/pot), a rating of 1 to 5 given, where 5 was complete desiccation of all plants

Two independent evaluations (ratings) done for plants treated at both 60 G/a and spray to drip

All treatments included 0.3% Hasten

Fatty acids at v/v and succinic acid (1%) and sodium salicylate (1%) at wt/v, in water

Caprylic/capric: caprylic, capric, caproic and lauric at 58, 40, 1 and 1% respectively (per Henkel Corporation)

No apparent correlation between pH of spray solutions (including Hasten) and herbicidal activity.

Example 25

Synergistic relationships between 5% caprylic/capric (c/c) and succinic acid: desiccation of cotton plants

Average Rating	Average percent of foliage desiccated	Treatment
2	20	
4	70	
4	60	0.5% succinic and
4	70	1% succinic and
		2% succinic and
		5% c/c
		5% c/c
		5% c/c
		5% c/c

* Average rating based on 1-5, where 5 = complete desiccation of foliage
Test plant: DeltaPine NuCotn 33-b, open boll
c/c = Caprylic, capric, caproic and lauric at 58, 40, 1 and 1%, respectively; i.e.,
Henkel's Emery 658, used at v/v, in water
All treatments contained 1% Emsorb 6900

Treatment of plants with succinic acid, alone, had a rating of zero
caprylic/capric and succinic at these concentrations were synergistic

Example 26

Synergistic relationships between 5% caprylic/capric (c/c) and succinic acid

<u>Treatment Ingredients</u>		
<u>Average rating: degree of plant damage</u> <u>(1-5, where 5 = complete desiccation)</u>		
<u>60 G/acre</u>	<u>Spray to drip</u>	
4.14	4.61	5% c/c
4.33	4.83	0.5% succinic and 5% c/c
4.43	4.82	1% succinic and 5% c/c
4.41	4.71	2% succinic and 5% c/c

Test plants: cotton (DeltaPine NuCotn 33b), potatoes (Snowden), pigweed, sudan grass, wheat, foxtail, dry beans and velvetleaf
c/c = Caprylic, capric, caproic and lauric at 58, 40, 1 and 1%, respectively; i.e., Henkel's Emery 658, used at v/v, in water

All treatments contained 1% Emsorb 6900

Treatment of plants with succinic acid, alone, had a rating of zero
Caprylic/capric (c/c) and succinic acid were synergistic

Example 27

Synergistic relationship between caprylic/capric (c/c) and succinic acid, as harvest aids for potatoes and cotton

Treatment Ingredients

Average rating: degree of plant damage
(1-5, where 5 = complete desiccation)

<u>Average, 2 evaluations</u>	
<u>Cotton</u>	<u>Potatoes</u>
3.75	4.1
4.35	4.25
4.65	4.15
4.65	4.35
	8% c/c
	0.5% succinic and 8% c/c
	1% succinic and 8% c/c
	2% succinic and 8% c/c

Cotton (DeltaPine NuCotn 33b) and potatoes (Snowden) were at full maturity
Application volume/acre was 40 G/a
C/C = Caprylic, capric, caproic and lauric at 58, 40, 1 and 1%, respectively; i.e.,
Henkel's Emery 658, used at v/v, in water
Treatment of plants with succinic acid, alone, had a rating of zero
Caprylic/capric (C/C) and succinic acid is synergistic

Example 28

Herbicidal activities: synergy between succinic acid and pelargonic acid

Treatment	Total injury ratings*			Spray to drip		Total
	40 G/a	60 G/a				
1 4% pelargonic acid	41.6	45.6		66.2		153.4
2 4% pelargonic acid + 0.5% succinic acid	42.8	46.7		66.3		155.8
3 0.5% succinic acid	0	0		0		0

*For 40 G/a: 9 total pots/treatment, testing crab grass, pigweed, chick weed and foxtail

*for 60 G/a: 10 total pots/treatment, testing pigweed, Kentucky bluegrass, foxtail, crab grass and barnyard grass

*For spray to drip: 7 total pots/treatment, testing Kentucky bluegrass, pigweed, barnyard grass, foxtail and velvetleaf

For each pot (2-20 plants/pot), a rating of 1-5 given, where 5 was complete desiccation of all plants. All treatments included 1.125% Emsorb 6900 (v/v)

Example 29

Synergistic relationships with caprylic/capric acids (c/c)

<u>Average relative injury rating*</u>		<u>Treatment Ingredients</u>	
<u>60 G/acre</u>	<u>Spray to drip</u>	<u>Average rating per pot</u>	<u>Compound (1%)</u>
4.5	4.57	4.52	urea
4.49	4.59	4.54	sorbic acid
4.56	4.33	4.45	di-sodium succinate
4.46	4.33	4.4	ammonium tartrate
4.62	4.5	4.58	methyl salicylate
4.53	4.52	4.53	potassium salicylate
4.56	4.47	4.52	succinic acid
4.6	4.61	4.61	

*For 60 G/a: 10 total pots/treatment, testing lambsquarter, pigweed, foxtail and barnyard grass
 *For spray to drip: 5 total pots/treatment, testing pigweed, sudan grass, foxtail, barnyard grass and velvetleaf

For each pot (2-20 plants/pot), a rating of 1 to 5 given, where 5 was complete injury/desiccation of all plants
 c/c = Caprylic, capric, caproic and lauric at 58, 40, 1 and 1%, respectively; i.e.,
 Henkel's Emery 658, used at v/v, in water

Example 30

Herbicidal activity on cotton foliage: synergistic relationships between succinic acid and caprylic/capric acid (c/c) combinations

Treatment		Total injury ratings, combined, for two separate evaluations*	
		60 G/a	Average rating/pot
1	4% c/c	18.5	3.08
2	4% c/c + 0.5% succinic acid	21.4	3.57
3	5% c/c	22.4	3.73
4	5% c/c + 0.5% succinic acid	25.7	4.28
5	6% c/c	25.3	4.22

*For 60 G/a: 3 total pots/treatment, testing cotton (DeltaPine NuCotn 33b)

For each pot, a rating of 1-5 was given, where 5 was complete desiccation of all foliage
Two independent evaluations (ratings) completed

All treatments included Emsorb 6900 (Henkel) at 1%

Caprylic/capric (caprylic, capric, caproic and lauric at 58, 40, 1 and 1%, respectively),
Henkel's Emery 658, used at v/v, in water

Since succinic acid tested alone had a rating of zero, caprylic/capric + succinic acid was synergistic
Herbicidal activity: 5% c/c + 0.5% succinic acid > 6% c/c

Example 31

Herbicidal activities: synergistic relationships between succinic acid and caprylic/capric acid (c/c) combinations

Treatment	Total injury ratings, combined for two separate evaluation*				Average rating per pot
	60 G/a	Spray to drip	Total Rating		
1 4% c/c	123.3	41.5	164.8		4.12
2 4% c/c + 0.5% succinic acid	128.1	45.1	173.2		4.33
3 5% c/c	129.9	46.3	176.2		4.41
4 5% c/c + 0.5% succinic acid	134.5	46.5	181		4.53
5 6% c/c	133.9	46.8	180.7		4.52

*For 60 G/a: 10 total pots/treatment, testing nightshade, pigweed, foxtail barnyard grass, sudan grass and cotton (DeltaPine NuCotn 33b)

*For spray to drip: 5 total pots/treatment, testing Kentucky bluegrass, nutsedge, barnyard grass, sudan grass and foxtail

For each pot (2-20 plants/pot), a rating of 1 to 5 given, where 5 was complete desiccation of all plants

Two independent evaluations (ratings) done for plants treated at both 60 G/a and spray to drip

All treatments included Emsoorb 6900 (Henkel) at 1%

Caprylic/capric (caprylic, capric, caproic and lauric at 58, 40, 1 and 1%, respectively), Henkel's Emery 658, used at v/v, in water
Since succinic acid, alone had a rating of zero, caprylic/capric + succinic acid was synergistic

General screening of candidate compounds: to examine for synergistic relationships with caprylic/capric acids (c/c)

*For 40 G/a: 10 total pots/treatment, testing chickweed, lambsquarter, Kentucky bluegrass, redtop and rye
 *For 60 G/a: 10 total pots/treatment, testing giant and green foxtail, pigweed and crabgrass
 *For spray to drip: 7 total pots/treatment, testing Kentucky bluegrass, nutsedge, pigweed, Johnson grass, foxtail and sudan grass
 For each pot (2-20 plants/pot), a rating of 1 to 5 given, where 5 was complete injury/desiccation of all plants
 c/c = Caprylic, capric, caproic and lauric at 58, 40, 1 and 1%, respectively; i.e., Henkel's Emery 658, used at v/v, in water.

Example 33

Herbicidal activity: storage stability of fatty acid/succinic acid-based formulations

Formulations stored one week (average day and night time temperatures were 90 and 75 degrees, Fahrenheit, in the greenhouse) and re-tested, via spray to drip applications on same test plant varieties

4% caprylic, 4% caprylic/capric and 4% pelargonic (+/- 0.5% succinic acid) were tested or a total of 6 formulations, each containing 1.125% Emsorb 6900 (Henkel)

Average scores, for all 6 formulations
(rating per pot, 2-20 plants/pot)

	<u>Before storage</u>	<u>After storage</u>
Barnyard grass	4.73	4.8
Foxtail	4.68	4.68

A rating of 1-5 was given, where 5 was complete desiccation of all plants

Fatty acids were v/v and succinic acid was wt/v, in water

Note: ratings for individual fatty acid treatments (+/- succinic acid), before and after storage, were similar

Example 34

Herbicidal combinations of caprylic/capric (c/c) or pelargonic acid, with equimolar amounts of succinic acid and ammonium succinate, on weed varieties

Treatment	Total Injury ratings*			Spray to drip	Total	Average rating/pot
	40 G/a	60 G/a	60 G/a			
1 3% c/c	85.5	60.2	60.2	36.1	181.8	4.33
2 3% c/c + 1% succinic acid	88.1	62.5	62.5	36.7	187.3	4.46
3 3% c/c + 1.26% ammonium succinate	88.8	60.8	60.8	37.4	187	4.45
4 3% pelargonic acid	87.7	60.8	60.8	37.3	185.8	4.42
5 3% pelargonic acid + 1% succinic acid	865.9	62.2	62.2	38	187.1	4.45
6 3% pelargonic acid + 1.26% ammonium succinate	87.6	61.2	61.2	36.9	185.7	4.42

*For 40 G/a: 10 total pots/treatment, testing crab grass, chickweed and common lambsquarter

*For 60 G/a: 7 total pots/treatment, testing redtop, pigweed, foxtail

For spray to drip: 8 total pots/treatment, testing barnyard grass, foxtail, sudan grass and nutsedge

For each pot (2-20 plants/pot), a rating of 1 to 5 given, where 5 was complete desiccation of all plants

Two independent evaluations for both 40 and 60 G/a and one evaluation for spray to drip

All treatments included Emsorb 6900 (Henkel) at 1%

Caprylic/capric (caprylic, capric, caproic and lauric at 58, 40, 1 and 1% respectively), i.e., Henkel's Emery 658, used at v/v, in water

Amounts of succinic acid and ammonium succinate used were equimolar amounts

Since succinic acid (or ammonium succinate) tested alone had a rating of zero, pelargonic acid (or c/c) + succinic acid (or ammonium succinate) were synergistic. The exception was: 3% pelargonic acid + 1.26% ammonium succinate

Example 35

Enhancement of RoundUp Ultra (+/- ammonium sulfate or Ams) herbicidal activity, using succinic acid amendments
RoundUp Ultra at 1 pint/acre (No Sylgard or any other surfactant used)

		<u>Days after treatment: relative injury ratings*</u>			<u>Average rating per pot</u>
		<u>Day #6</u>	<u>Day #9</u>	<u>Day #11</u>	
1	RU	10.5	17.1	15.9	1.91
2	RU + ams	18.3	17.6	18.4	2.37
3	RU + ams + 0.5% succinic acid	21	20.3	24.3	2.85
4	RU + ams + 1.5% succinic acid	19.4	18.6	24.2	2.8
5	RU + 0.5% succinic acid	16.7	14.2	16	2.11
6	RU + 1.5% succinic acid	16.8	16.1	18.6	2.35

*8 total pots/treatment, testing Kentucky bluegrass, barnyard grass, crab grass, pigweed, redtop, nutsedge, wheat and foxtail. For each pot (2-20 plants/pot), a rating of 1 to 5 given, where 5 was complete injury/desiccation of all plants
 RoundUp Ultra was at 1 pint/acre and ammonium sulfate, when used, was at 2%
 Spray application at 60 G/acre
 Succinic acid tested alone had a rating of zero. RoundUp Ultra, alone or RoundUp Ultra + ammonia sulfate were synergistic with succinic acid

Example 36

Enhancement of RoundUp Ultra (+/- ammonium sulfate or Ams) herbicidal activity, using succinic acid amendments
RoundUp Ultra at 1 pint/acre

Treatment	Days after treatment: relative injury ratings*				Average rating per pot
	Day #6	Day #9	Day #11	Day #14	
1 RU	29	30.2	31.5	29.8	3.35
2 RU+ams	30.8	32.9	34.7	32.5	3.64
3 RU+ams + 0.5% succinic acid	30.2	32.7	35.5	34.5	3.69
4 RU+ams + 1.5% succinic acid	34.6	35.7	37.8	37.4	4.04
5 RU + 0.5% succinic acid	32.8	33.9	35.1	35.4	3.81
6 RU + 1.5% succinic acid	32.5	31.2	32.6	32.5	3.58

*9 total pots/treatment, testing barnyard grass, green foxtail, sudan grass and pigweed. For each pot (2-20 plants/pot), a rating of 1 to 5 given, where 5 was complete injury/desiccation of all plants. RoundUp Ultra was at 1 pint/acre and ammonium sulfate, when used, was at 2% Sylgard 309 (0.3%) added immediately before spray application at 60 gallons/acre. Succinic acid tested alone had a rating of zero. RoundUp Ultra, alone or RoundUp Ultra + ammonium sulfate were synergistic with succinic acid.

Example 37

Enhancement of RoundUp Ultra (+/- ammonium sulfate or Ams) herbicidal activity, using succinic acid amendments
RoundUp Ultra (RU) at 1 quart per acre

Treatment	Days after treatment: relative injury ratings*					Average rating per pot
	Day #5	Day #8	Day #10	Day #13	Day #17	
1 RU	32.7	29.9	28.6	29.5	31.4	3.8
2 RU+ams	33.3	33.6	30.7	34.5	36.6	4.22
3 RU+ams + 0.5% succinic acid	34	34.5	32.4	35.3	36.7	4.32
4 RU+ams + 1.5% succinic acid	35.3	35.6	32.5	35.3	38.4	4.43
5 RU + 0.5% succinic acid	33.8	30.9	30.3	31	33.1	3.98
6 RU + 1.5% succinic acid	33.7	30.2	31	33.6	34	4.06

*8 total pots/treatment, testing sudan grass, green foxtail, barnyard grass and pigweed
 For each pot (2-20 plants/pot), a rating of 1 to 5 given, where 5 was complete injury/desiccation of all plants
 RoundUp Ultra was at 1 quart/acre and ammonium sulfate, when used, was at 2%, wt/v
 Sylgard 309 (0.3%) added immediately before spray application at 60 gallons/acre
 Succinic acid tested alone had a rating of zero. RoundUp Ultra, alone or RoundUp Ultra + ammonium sulfate were synergistic with succinic acid

Example 38

Enhancement of herbicidal activity of RoundUp Ultra/ammonium sulfate testing selected amendments

Treatment	pH, spray solution	Days after treatment: injury ratings*				average per test
		day #4	day #7	day #11	day #13	
1 RU/ams	4.51	32.3	44.1	50	50.8	4.03
2 RU/ams+succinic acid	1.77	39.3	46.3	50.7	51.4	4.26
3 RU+ams+ammonium succinate	3.71	36.2	46.2	51.3	52.3	4.23
4 RU+ams ammonium tartrate	4.78	35.7	46	49.3	50.2	4.12
5 RU/ams+tartaric acid	1.27	33.9	46.9	50.7	51.5	4.2
6 RU/ams+citric acid	1.38	34.6	46.7	50.7	51.4	4.15
7 RU/ams+L-malic acid	1.59	34.6	47.7	51.1	51.8	4.21
8 RU/ams+acetic acid	2.51	33.7	47.1	50.8	51.3	4.15

*11 total pots/treatment, testing velvetleaf, crab grass, sudan grass, redtop, green foxtail and barnyard grass. For each pot (2-20 plants/pot), a rating of 1 to 5 given, where 5 was complete desiccation

Equimolar amounts of amendments used: 1.5% succinic acid, 1.89% ammonium succinate, 2.17% ammonium tartrate, 1.91% tartaric acid, 2.45% citric acid, 1.71% L-malic acid, 0.77% acetic acid

RoundUp Ultra (1.5 pints/acre) and 2% ammonium sulfate (ams) used in all treatments
Sylgard (0.3%) added immediately before spray application at 60 gallons/acre

Amendments used independently had a rating of zero. All amendments had a synergistic relationship with RU/ams; i.e., each amendment enhanced RU/ams herbicidal activity over ratings for RU/ams (containing no amendments). No apparent correlation between pH of spray solutions (including Sylgard) and herbicidal activity was found.

Example 39

Enhancement of herbicidal activity of caprylic/capric acids (c/c), using selected compounds at equimolar amounts

Treatment	Days after treatment:				Average rating per pot
	Day #2	Day #4	Day #5	Day #5	
1 3% c/c	30	27.6	30.3	30.3	3.66
2 3% c/c + succinic acid (1%)	35.8	33.6	35.1	35.1	4.35
3 3% c/c + di-ammonium succinate (1.26%)	35.1	32.5	34.2	34.2	4.24
4 3% c/c + ammonium sulfate (1.12%)	35.5	33.6	34.4	34.4	4.3
5 3% c/c + ammonium tartrate (1.42%)	35.9	33.6	34.2	34.2	4.32
6 succinic acid (1%)	0	0	0	0	0
7 di-ammonium succinate (1.26%)	0	0	0	0	0
8 ammonium sulfate (1.12%)	0	0	0	0	0
9 ammonium tartrate (1.42%)	0	0	0	0	0

*8 total pots/treatment, testing chickweed, redtop, velvetleaf, crabgrass and green foxtail
 For each pot (2-20 plants/pot), a rating of 1 to 5 given, where 5 was complete injury/desiccation of all plants
 All treatments included 0.86% Emsorb 6900 and 0.43% mineral oil
 c/c = Caprylic, capric, caproic and lauric at 58, 40, 1 and 1% respectively; i.e.,
 Henkel's Emery 658, used at v/v, in water
 Application volume was 40 G/acre

Synergistic relationships exist between 3% c/c and each amendment tested

Example 40

Desiccation of cotton foliage: Interaction of RoundUp Ultra and caprylic acid/capric acid (+/- succinic acid)

Treatment	Day after treatment: average rating*										Total Rating
	1	2	3	4	5	6	7	8	9	10	
1 RU + 2% AMS	0	0	0	0	0	0	0	0	0	0	26
2 RU + 2% AMS	18	16	15	15	15	15	15	15	15	15	95.9
3 RU + 2% AMS	18	17	17	17	17	17	17	17	17	17	103.2
4 RU + 2% AMS	0	0	0	0	0	0	0	0	0	0	30.2
5 2% AMS	18	17	18	18	18	18	18	18	18	18	105.4
6 2% AMS	18	17	18	18	18	18	18	18	18	18	104.8

*Visual rating of 1 to 5, where 5= complete desiccation of all cotton plants (4 plants per treatment group, each plant receiving a rating of 1-5)

Cotton variety = DeltaPine NuCotn 33b, at open boll, when treated

RoundUp Ultra at 1 quart/acre delivered at 60 G/a

Each treatment contained 1.14% Emsorb 6900 and 0.57% mineral oil

Caprylic/capric (caprylic, capric, caproic and lauric at 58, 40, 1 and 1% respectively), i.e., Henkels, Emery 658, used at v/v, in water

AMS = ammonium sulfate

Succinic acid had a rating of zero.

Combination of succinic acid with other ingredients was synergistic.

Example 41

Synergistic relationship between succinic acid and caprylic/capric acids C/C, relating to herbicidal activity

Injury ratings based on 2 independent evaluations*

<u>Treatment</u>	<u>40 G/a</u>	<u>60 G/a</u>	<u>Total</u>	<u>Average rating per pot</u>
1 4% caprylic/capric	87.3	80.7	168	4.42
2 4% caprylic/capric + 1% succinic	88.6	82.1	170.7	4.49
3 4% caprylic/capric + 0.5% succinic	89.9	82.7	172.6	4.54
4 4% caprylic/capric + 0.5% succinic + 0.5% sodium salicylate	90.3	82.6	172.9	4.55
5 0.5% succinic acid	0	0	0	0
6 0.5% sodium salicylate	1	0.5	1.5	0.04
7 5% caprylic/capric	92	82.8	174.8	4.6
8 5% caprylic/capric + 0.5% succinic acid	91.6	83.9	175.5	4.62

*For each pot, a rating (1 to 5) was given, where 5 was complete desiccation of all plants

Two independent evaluations were made for both 40 and 60 gallons/acre applications

For 40 G/acre: Canadian thistle, velvet leaf, Johnson grass and giant foxtail

For 60 G/acre: velvetleaf, Johnson grass and giant foxtail were tested

C/C = Caprylic, capric, caproic and lauric at 58, 40, 1 and 1%, respectively; i.e.,

Henkel's Emery 658, used at v/v, in water

Succinic acid and sodium salicylate were added at wt/volume

All treatments included Henkel's Emery 6900 and mineral oil; i.e., for each 1% of caprylic/capric, 0.286 and 0.143% of 6900 and mineral oil were included in each treatment

Example 42

Herbicidal activity: comparison of caprylic, pelargonic and an equimolar mixture of caprylic and pelargonic acids

<u>Injury ratings based on 2 independent evaluations*</u>			<u>Average rating per pot</u>
<u>Treatments (equimolar comparison)</u>	<u>40 G/acre</u>	<u>60 G/acre</u>	
1 3% caprylic	40.8	51.6	4.16
2 3.375% pelargonic	43	53.6	4.41
3 1.5% caprylic + 1.69% pelargonic	43	53.6	4.41
4 4% caprylic	44.1	54.5	4.49
5 4.5% pelargonic	45.1	53.9	4.51
6 2% caprylic + 2.25% pelargonic	44.6	54.2	4.51

*For each pot, a rating (1 to 5) was given, where 5 was complete desiccation of all plants
 For 40 G/acre: chickweed, redroot pigweed, lambsquarter and green foxtail were tested
 For 60 G/acre: chickweed, redroot pigweed, lambsquarter, green foxtail, crabgrass and wheat were tested
 Caprylic and pelargonic acids were used at volume/volume
 All treatments included Henkel's Emsorb 6900 and mineral oil: for each 1% of caprylic or pelargonic acid, 0.286 and 0.143% of 6900 and mineral oil were included in each treatment.

Example 43

Herbicidal enhancement of pelargonic by succinic acid and diammonium succinate

<u>Injury ratings based on 2 independent evaluations*</u>					
<u>Treatment</u>	<u>40 G/a</u>	<u>60 G/a</u>	<u>Spray to drip</u>	<u>Average rating per pot</u>	
1 2.5% Scythe active ingredients	78.7	103	53.6	4.36	
2 2.5% pelargonic	79.2	102.2	53.8	4.36	
3 2.5% pelargonic + 0.5% succinic	81.7	105.1	55.3	4.48	
4 2.5% pelargonic + 2% succinic	80.2	105.5	55.1	4.46	
5 2.5% pelargonic + 2% diammonium succinate	80.9	105.4	54.4	4.46	
6 2.22% caprylic/capric (C/C)	80.3	103.4	51.5	4.36	

*For each pot, a rating (1 to 5) was given, where 5 was complete desiccation of all plants

40 G/a: lambsquarter, velvetleaf and green foxtail were tested

60 G/a: lambsquarter, velvetleaf and green foxtail, velvetleaf, lambsquarter and wheat were tested

Spray to drip: nutsedge, velvetleaf, green foxtail, giant foxtail and barnyard grass were tested

C/C = Caprylic, capric, caproic and lauric at 58, 40, 1 and 1%, respectively, i.e.,

Henkel's Emery 658, used at v/v, in water

The active ingredients in Scythe, (pelargonic acid) and caprylic/capric (Henkel's Emery 658) were compared on equimolar basis

Example 44

Synergistic Relationship between Succinic or Citric acid and Caprylic Acid, Testing Dry Beans
Treatment effects 3 days after single, foliar application

Treatment	<u>Green Foliage</u>		<u>Yellow Foliage</u>	
	Overall Effect	Foliage Affected (%)	Overall Effect	Foliage Affected (%)
Na Salicylate (2%)	5	90	5	>90
Na Salicylate (0.5%)	2	5-10	1	30
Na Salicylate (0.5%) + Citric Acid (0.5%)	1	5-10	1	30
Na Salicylate (0.5%) + Succinic Acid (0.5%)	1.5	5-10	1	>40
Citric Acid (0.5%)	0	0	0	0
Succinic Acid (0.5%)	0	0	0	0
Caprylic Acid (0.5%)	2.5	10-15	2	70
Caprylic Acid (0.5%) + Citric Acid (0.5%)	3.5	30	3	90
Caprylic Acid (0.5%) + Succinic Acid (0.5%)	4	50	4	>90

Overall effect: higher scores = greater effect
 (where 5 equals profound desiccation)
 Dry bean variety = Vista

Citric acid and succinic acid showed synergistic effect

Example 45

Greater Effect by Succinic Acid over Calcium Succinate, when used with Sodium Salicylate, Testing Several Plant Varieties*

Treatments	overall
	effect, 9 evaluations (average)
Sodium Salicylate (1%) + Succinic Acid (1%)	2.89
Sodium Salicylate (1%) + Calcium Succinate (1%)	1.44

Overall effect: higher score = greater effect
(where 5 equals profound desiccation)

*Test plants: weed varieties, shrub foliage, covergrass, turf and soybeans

Example 46

Confirmation of Example 45, Testing Turf and Shrub Foliage
Treatment effects 1, 2, 4 and 7 days after a single, foliar application.

Treatments	Overall effect, 17 evaluations <u>(average)</u>
Sodium Salicylate (1%)	1.35
Sodium Salicylate (1%) + Succinic Acid (2%)	2.59
Sodium Salicylate (1%) + Calcium Succinate (2%)	2.18
Succinic Acid (2%)	0.24

Overall effect: higher score = greater effect
(where 5 equals profound desiccation)

Succinic acid and calcium succinate showed synergistic effect

Example 47

Effects of desiccants and desiccant combinations with other compounds on young potato plants

Treatment #	Ingredients	Injury ratings of treated plants (hours or days after application)						All evaluations (total score)
		1.5 hr	1	2	3	6	17	
1	1% caprylic	2	2	2	3	2	2.5	27
2	1% caprylic + 0.5% tartaric + 0.5% NaSal	3	5	5	5.5	5	5.5	3
3	1% caprylic + 1% tartaric	2	3	3	3.5	3	3.4	4
4	1% caprylic + 0.5% dimethyl succinic acid	2	3	3	3	3	3.5	3
5	1% caprylic + 0.5% beta-alanine	1	2	2	2	2	2.5	3
6	1% caprylic + 1% succinic	3	4	4	3	4	5	2.5
7	2% caprylic	3.5	5	5	5	5	5.5	4.5
8	2% caprylic + 0.5% tartaric + 0.5% NaSal	4.5	6	6	6	6	6	5.5
9	sodium salicylate	3.5	5	5	5	5	5	6
	2% caprylic + 1% tartaric							34.5
								40.5
								34.5

Caprylic acid and dimethyl succinic acid added v/v; all other compounds added at wt/v. All solutions in 50% acetone, with solution (20 mls) sprayed 2 feet from plant canopy. Helena Kinetic at 0.2% included in each solution. All plants grown in green house (day time temp= 80-100 degrees, Fahrenheit) Higher evaluation scores represent greater degree of vine and foliage desiccation. A "6" = complete desiccation. Summary: a) addition of succinic or tartaric acid to 1% caprylic acid>>1% caprylic acid, (b) combination of NaSal + tartaric (or succinic acid) to caprylic acid were the best treatments

Example 48

Synergistic Relationships Between:

Succinic Acid + Caprylic Acid

Succinic Acid + Sodium Salicylate

Treatment effects 3 days after a single, foliar application, testing soybeans

<u>Treatment</u>	<u>Overall Effect</u>	<u>Foliage Affected (%)</u>
Caprylic Acid (1%)	2.5	25-30
Caprylic Acid (1%) + Succinic Acid (1%)	3	30-35
Succinic Acid (1%)	0	0
Sodium Salicylate (1%)	1.5	15-20
Sodium Salicylate (1%) + Succinic Acid (1%)	3.5	40-50

Overall effect: higher scores = greater effect
(where 5 equals profound desiccation
Soybean variety = Stine 2250

Example 49

Enhancement of Scythe's herbicidal activity by selected compounds

Treatment #	Treatment Ingredients	Independent evaluations of snapbean and potato plants, hours/days after application of desiccants									
		1.5 hours		2 days		11 days		15 days		potatoes	
		beans	potatoes	beans	potatoes	beans	potatoes	beans	potatoes	beans	potatoes
1	2% NaSalicylate	1	0	1	1	1.5	1	1	1	1	potatoes
2	4% Scythe	3	3	3.5	2.5	2	2	1	2	2	2
3	4% Scythe + 2% NaSal	4	3	5	3.5	3	3	2	3	3	3
4	4% Scythe + 0.25% tartaric	4	3	4	3	2.5	2.5	3	2	2	2
5	4% Scythe + 0.5% tartaric	4	4	5.5	3	3	2	4	1.5	1.5	1.5
6	4% Scythe + 1% tartaric	4	3.5	6	3.5	3	3	3	2	4	1.5
7	4% Scythe + 1% NaSal	3	3	4.5	3	2.5	2.5	2	2	3	2
8	4% Scythe + 0.5% NaSal	3	3	4.5	2	1.5	1.5	1	1	1	2
9	4% Scythe + 1% caprylic	4	4.5	5	3.5	2	3.5	1	3.5	3	3.5
10	4% Scythe + 0.5% tartaric + 0.5% NaSal	4	5	5.5	5	3.5	4	3	5	3	5
11	4% Scythe + 1% citric acid	4	3	6	3	2.5	2	2	2	2	2

Totals, all evaluations			All
Beans	Potatoes		
1 4.5	3	7.5	
2 9.5	9.5	19	
3 14	12.5	26.5	
4 13.5	10.5	24	
5 16.5	10.5	27	
6 16	12	28	
7 12	10	22	
8 10	8	18	
9 12	15	27	
10 16	19	35	
11 14.5	10	24.5	

Solutions applied at 40 gallons/acre, containing 0.1% Helena Kinetic. All plants grown in greenhouse (day temperature was 80-100 degrees, fahrenheit). Higher rating scores represent greater degree of desiccation. A "6" = complete desiccation. Scythe and caprylic acid added v/v. All other compounds added w/v.

Summary: a) all combinations with Scythe (except 0.5% NaSal) improved performance, b) best combination was 4% Scythe with 0.5% tartaric + 0.5% NaSal, c) increasing the concentration of tartaric acid with 4% Scythe had little effect

Example 50

Synergistic Relationships Between Succinic Acid and Caprylic Acid at Different Application Rates, Testing Soybeans
Treatment effects 2 days after a single foliar application

	Overall Effect	Two Independent Sites (average of 2 sites)	Foliage Affected (%)
Caprylic Acid (0.2%)	0.5		1.25
Caprylic Acid (0.5%)	2.25		9
Caprylic Acid (1.0%)	4		22.3
Caprylic Acid (0.2%) + Succinic Acid (1.0%)	1.5		6.5
Caprylic Acid (0.5%) + Succinic Acid (1.0%)	3		15
Caprylic Acid (1.0%) + Succinic Acid (1.0%)	4.25		26
Succinic Acid (1.0%)	0		0
Caprylic Acid (0.5%) + L - Lactic Acid (1.0%)	2.75		15
L-Lactic Acid (1.0%)	0.5		1.25

Overall effect: higher scores = greater effect
(where 5 equals profound desiccation)
Soybean variety = Stine 2250

Succinic acid and lactic acid had synergistic effect

Example 51

Synergy Comparisons of Various Organic Acids with Caprylic Acid, Testing Soybeans

Treatment effects 3 days after a single, foliar application

Treatments	Two Independent Sites (average of 2 sites)	
	Overall Effect	Foliage Affected (%)
Caprylic Acid (0.5%)	1.5	5
Caprylic Acid (0.5%) + L-Tartaric Acid (0.5%)	2.75	17.5
Caprylic Acid (0.5%) + L-Malic Acid (0.5%)	2	11.5
Caprylic Acid (0.5%) + Succinic Acid (0.5%)	2.25	12.5
Caprylic Acid (0.5%) + L-Lactic Acid (0.5%)	2.25	11.5
Caprylic Acid (0.5%) + Citric Acid (0.5%)	2.75	17.5
L-Tartaric Acid (0.5%)	0	0
L-Malic Acid (0.5%)	0	0
Succinic Acid (0.5%)	0	0
L-Lactic Acid (0.5%)	0	0
Citric Acid (0.5%)	0	0

Overall effect: higher scores = greater effect
(where 5 equals profound desiccation)

Soybean variety = Stine 2250

Tartaric, malic, succinic, lactic and citric acid showed synergistic effects

Example 52

Synergistic Relationships Between Succinic Acid and Caprylic Acid (or Sodium Salicylate), Testing Turf

Treatment effects 2 days after a single, foliar application

Treatments	Overall Effect
Caprylic Acid (1.0%)	2
Caprylic Acid (1.0%) + Succinic Acid (0.5%)	3
Succinic Acid (0.5)	0
Sodium Salicylate (1.0%)	1.5
Sodium Salicylate (1.0%) + Succinic Acid (0.5%)	2

Overall effect: higher scores = greater effect
(where 5 equals profound desiccation)

Example 53

Effects of Various Compounds on Caprylic Acid Testing Turf and Covergrass

Treatment effects 1, 3, 6, 7 and 11 days after a single,
foliar application at two, independent sites

Treatments	Overall Effect 10 Evaluations (average)
Caprylic Acid (1.0%)	2.45
Caprylic Acid (1.0%) + Adipic Acid (0.5%)	2.6
Caprylic Acid (1.0%) + L-Tartaric Acid (0.5%)	3.8
Caprylic Acid (1.0%) + Unipine (0.5%)	3.45
Caprylic Acid (1.0%) + Sodium Salicylate(0.5%)	3.1

Overall effect: higher scores = greater effect
(where 5 equals profound desiccation)

Adipic acid, tartaric acid, unipine and sodium salicylate showed synergistic effects

Example 54

Effects of Various Compounds on Caprylic Acid Testing Turf and Shrub Foliage

Treatment effects 3, 4, 6 and 7 days after a single, foliar application

Treatments	Overall Effect 29 Evaluations (average)
Caprylic Acid (1.0%)	1.05
Caprylic Acid (1.0%) + Adipic Acid (0.5%)	2.19
Caprylic Acid (1.0%) + L-Malic Acid (0.5%)	1.4
Caprylic Acid (1.0%) + L-Tartaric Acid (0.5%)	2.19
Caprylic Acid (1.0%) + Unipine (0.5%)	2.4
Caprylic Acid (1.0%) + Sodium Salicylate(0.5%)	2.05
Caprylic Acid (1.0%) + Succinic Acid (0.5%)	1.95
Caprylic Acid (1.0%) + L-Lactic Acid (0.5%)	2.16
Caprylic Acid (1.0%) + Citric Acid (0.5%)	1.47

Overall effect: higher scores = greater effect
(where 5 equals profound desiccation)

Adipic acid, tartaric acid, unipine, sodium salicylate, succinic acid, lactic acid and citric acid showed synergistic effects

Example 55

Measurement of pH Values for Test Desiccants

Caprylic Acid (0.5%)	pH Value
Caprylic Acid (0.5%) + L-Tartaric Acid (0.5%)	3.30
Caprylic Acid (0.5%) + L-Malic Acid (0.5%)	1.07
Caprylic Acid (0.5%) + Succinic Acid (0.5%)	1.49
Caprylic Acid (0.5%) + L-Lactic Acid (0.5%)	1.67
Caprylic Acid (0.5%) + Citric Acid (0.5%)	1.69
L-Tartaric Acid (0.5%)	1.51
L-Malic Acid (0.5%)	1.52
Succinic Acid (0.5%)	1.69
L-Lactic Acid (0.5%)	1.94
Citric Acid (0.5%)	1.93
	1.64
Caprylic Acid (0.5%)	3.55
Caprylic Acid (0.5%) + Succinic Acid (0.5%)	1.77
Succinic Acid (1.0%)	1.85
Sodium Salicylate (1.0%)	5.63
Sodium Salicylate (1.0%) + Succinic Acid (0.5%)	2.9
Caprylic Acid (1.0%)	3.93
Caprylic Acid (1.0%) + Adipic Acid (0.5%)	2.69
Caprylic Acid (1.0%) + L-Malic Acid (0.5%)	2.17
Caprylic Acid (1.0%) + L-Tartaric Acid (0.5%)	2.07
Caprylic Acid (1.0%) + Unipine 90 (0.5%)	4.06
Caprylic Acid (1.0%) + Sodium Salicylate (0.5%)	4.29

Example 56

Measurement of pH Values for Test Desiccants

	pH Value
Sodium Salicylate (0.5%)	4.48
Sodium Salicylate (0.5%) + Succinic Acid (1.0%)	2.76
Sodium Salicylate (0.5%) + Succinic Acid (1.0%) + Caprylic Acid (0.5%)	2.78
Sodium Salicylate (0.5%) + Caprylic Acid (0.5%)	4.2
Succinic Acid (1.0%) + Caprylic Acid (0.5%)	2.24
Sodium Salicylate (0.5%) + Caprylic Acid (0.5%) + L-Tartaric Acid (1.0%)	2.46
Sodium Salicylate (0.5%) + Caprylic Acid (0.5%) + L-Lactic Acid (1.0%)	2.82
Sodium Salicylate (0.5%) + Caprylic Acid (0.5%) + Succinic Acid (1.0%)	3.24
Sodium Salicylate (0.5%) + Caprylic Acid (0.5%) + Succinic Acid (1.0%)	2.32
Sodium Salicylate (0.5%) + Caprylic Acid (0.5%) + L-Lactic Acid (1.0%)	2.02
Sodium Salicylate (0.5%) + Caprylic Acid (0.5%) + L-Tartaric Acid (1.0%)	1.58
Sodium Salicylate (0.5%) + Caprylic Acid (0.5%) + Unipine 90 Acid (1.0%)	5.12
Sodium Salicylate (0.5%) + Caprylic Acid (0.5%) + Adipic Acid (1.0%)	2.78
Caprylic Acid (1.0%) + Gluconic Acid (0.5%)	2.49
Caprylic Acid (1.0%) + Succinic Acid (0.5%)	2.29
Caprylic Acid (1.0%) + Beta-Alanine (0.5%)	4.4
Caprylic Acid (1.0%)	3.85
Caprylic Acid (1.0%) + L-Tartaric Acid (0.5%)	1.76
Caprylic Acid (1.0%) + L-Lactic Acid (0.5%)	1.95
L-Tartaric Acid (0.5%)	1.72
Beta-Alanine (0.5%)	6.65

Example 57

Field trial comparisons of caprylic/capric and pelargonic acids on mixed weed varieties

<u>Treatment</u>	<u>Average injury rating per plot*</u>
1 3% caprylic/capric	3.95
2 3% caprylic/capric + 0.75% succinic acid	4.113
3 3% pelargonic	4.175
4 3% pelargonic + 0.75% succinic acid	4.213

* A rating of 1-5 given for each of 2 plots per treatment group where 5 was complete desiccation of all plants

Plot size: 6 X 9.3 feet

Weed varieties: grasses (dominant), dandelion, pigweed, plantain and Canadian thistle

Summary: addition of succinic acid was beneficial

Example 58

Field trial comparisons of caprylic/capric and pelargonic acids on mixed weed varieties

Canadian thistle:		Average injury rating per plot*
Treatment		
1	3% caprylic/capric	2.58
2	3% caprylic/capric + 0.75% succinic acid	2.8
3	3% pelargonic	4.03
4	3% pelargonic + 0.75% succinic acid	4.35

Redwood pigweed:		Average injury rating per plot*
Treatment		
1	4% caprylic/capric	3.1
2	6% caprylic/capric	4.55
3	6% pelargonic	4.35

* A rating of 1-5 given for each of 2 plots per treatment group where 5 was complete desiccation of all plants
Plot size: 6 X 9.3 feet

Summary: 1) although pelargonic acid was more effective for C. Thistle, caprylic/capric performed best on redroot pigweed, 2) addition of succinic acid was effective

Example 59

**Succinic Acids Potentiation of Sodium Salicylate,
Testing Cotton**

Treatment effect 2 days after a single, foliar application

<u>Treatment</u>	<u>Foliage</u>	
	<u>Affected (%)</u>	<u>No effect</u>
Succinic Acid (1.0%)		
Succinic Acid (1.0%) + Sodium Salicylate (1.0%)	>70	
Sodium Salicylate (1.0%)	<70	

Some foliage (all treatments except succinic acid, alone) beginning to drop

It will thus be seen that the objects set forth above, among those made apparent from the preceding description, are efficiently attained and, since certain changes may be made in carrying out the above methods and in the compositions set forth, without departing from the spirit and scope of the invention, it is intended that all matter contained in the above description shall be interpreted as illustrative and not in a limiting sense.

It is also understood that the following claims are intended to cover all of the generic and specific features of the invention herein described and all statements of the scope of the invention which, as a matter of language might be said to fall therebetween.

Particularly, it is to be understood that in said claims, ingredients or components recited in the singular are intended to include compatible mixtures of said ingredients wherever the sense permits.